Cleans, sanitizes<sup>3</sup>, disinfects<sup>2</sup> and deodorizes. All Purpose Cleaner. An effective soft surface sanitizer § as a spot treatment. Designed for use in refillable spray bottle, mechanical spray device, low pressure sprayer and trigger spray. Refer to equipment manual for operation instruction before use. This product will clean and disinfect<sup>2</sup> hard, non-porous surfaces of: grocery cart handles, cribs, diaper pails, trashcans, garbage cans, gym equipment, exercise machines, wrestling mats, avmnastic equipment.

#### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not for use on critical and semi-critical medical device surfaces. Fill or refill the spray bottle with this product. Pour product into spray bottle over sink or tub basin. GENERAL CLEANING: Spray soiled area then wipe with a dry paper towel or lint-free cloth. DISINFECTING, DEODORIZING: Remove visible fillth or soil prior to application of the product. Apply product with a mechanical spray device, cord-less backpack sprayer, trigger spray or hand held mechanical sprayer. Hold sprayer six to eight inches from the surface to be treated. Spray area until it is covered with the product. Allow surface to remain visibly wet for 2 minutes. No scrubbing necessary. Allow treated surfaces to air dry or wipe off with a clean cloth. When using on food contact surfaces: counter tops, room-temperature stovetops, highchairs, kitchen appliances, thoroughly rinse all treated surfaces with potable tap water. This product must not be used to clean

eating utensils, glassware and dishes.

This product is an effective disinfectant against: Pseudomonas aeruginosa, Salmonella enterica, Staphylococcus aureus, Methicillin Resistant Staphylococcus aureus, Vancomycin Resistant Enterococcus faecium, Multi-Drug Resistant Acinetobacter baumanni Ceftazidime and Gentamicin Resistant, Extended Spectrum Beta-Lactamase Escherichia coli and Klebsiella pneumoniae New Delhi Metallo-Beta Lactamase Carbapenem Resistant.

\*VIRUCIDAL ACTIVITY: This product is an effective virucide on hard, non-porous surfaces against Human Immunodeficiency Virus Type-1(HIV-1) and Avian Influenza Virus NIBRG-14, and SARS-Related Coronavirus 2 (SARS-CoV-2) when the treated surface is allowed to remain visibly wet for 30 seconds. This product is an effective virucide on hard, non-porous surfaces against Human Rotavirus, Hepatitis B Virus(HBV), Hepatitis C Virus(HCV), Duck Hepatitis B Virus(BHSV) and Bovine Viral Diarrhea Virus (BVDV) when the treated surface is allowed to remain visibly wet for 1 minute. This product is an effective virucide on hard non-porous surfaces against Feline Calicivirus, Norovirus (Norvalk Virus), Canine Parvovirus, Rhinovirus Type 14 and Rhinovirus Type 39, when the treated surface is allowed to remain visibly wet for 2 minutes. This product is effective against Poliovirus Type 1 when the treated surface is allowed to remain visibly wet for 5 minutes. Allow treated surfaces to air dry or wipe off with a clean cloth.

KILLS HIV-1, HBV and HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1, Hepatitis B Virus and Hepatitis C Virus.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV AND HCV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

PERSONAL PROTECTION: When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks, and eye coverings, CLEANING PROCEDURES: Blood



# QUICK CLEAN RTU Disinfectant

Kills 99.9%<sup>2</sup> of SARS-CoV-2, which causes COVID-19<sup>5</sup> on hard non-porous surfaces in 30 seconds Cleans, Disinfects<sup>2</sup> & Deodorizes. \*Virucidal

## KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. If IN EYES: Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

See side panel for additional precautionary statements.

Net Contents: 1 Quart U.S.

631012 01/22

and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **CONTACT TIME:** Allow surfaces to remain visibly wet for 2 minutes. HIV-1 is inactivated in 30 seconds, HBV and HCV are inactivated in 1 minute.

DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

SANITIZE NON-FOOD CONTACT SURFACES and DEODORIZE – To sanitize/deodorize hard, non-porous surfaces, hold container six to eight inches from the surfaces to be treated. Spray area until it is covered with the product. Allow surface to remain visibly wet for 5 seconds. This product is an effective sanitizer against Staphylococcus aureus (Staph) and Klebsiella pneumoniae.Let surface dry. For visibly soiled surfaces, a precleaning step is required.

SOFT SURFACE SÁNITIZATION – To Sanitize Soft Surfaces (Fabrics): Hold bottle upright 6"-8" from surface. Spray until fabric is wet. DO NOT SATURATE. Surface must remain visibly wet for 2-minute contact time. Allow to air dry. Reapply as necessary. Visible soil must be removed prior to sanitizing. This product is an effective sanitizer against Klebsiella aerogenes and Staphylococcus aureus in 2 minutes. For first time use on soft surfaces, spot test in a small, inconspicuous area.

#### STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

PESTICIDE STORAGE - Store in a dry place inaccessible to children.

CONTAINER HANDLING – Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or puncture and dispose in a sanitary landfill, incineration, or burning, if allowed by state and local authorities. If burned, stay out of smoke.

CONTAINER DISPOSAL FOR REFILL SPRAY BOTTLE: – Do not reuse empty spray bottle except with refill container, otherwise wrap spray bottle and discard in trash. Offer for recycling if available. PESTICIDE DISPOSAL – Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

- composed of 100% cotton or 100% polyester or 50/50 cotton polyester.
- 2 When used according to disinfection directions 5 SARS-Related Coronavirus 2
- 3 When used according to hard surface sanitization directions

SUNBELT LABORATORIES
P.O. BOX 1563, STAFFORD, TX 77497-1563
281-261-4747/ 1-800-324-4413 • www.sunbelt-labs.com